

Improving Morbidity and Mortality Associated with Chronic Kidney Disease

Zeidan Hammad, MD, FAAFP

Assistant Professor of Family Medicine
Department of Humanities, Health, & Society
Florida International University
Herbert Wertheim College of Medicine
Florida, USA

Elaine Huang

Medical Student
Herbert Wertheim College of Medicine
Florida, USA


Abstract

The rising prevalence of chronic kidney disease raises questions on the approaches that can be taken to decrease morbidity, mortality, and cost. Some measures that have been discussed suggest adequate control of chronic kidney disease (CKD) risk factors, such as diabetes and hypertension. Potential avenues of mortality and cost reduction such as early screening, a multidisciplinary model of care, and primary care involvement are discussed here.

Introduction

The incidence of chronic kidney disease in the United States is one of the highest in the world, “30 million people or 15% of US adults are estimated to have CKD, 48% of those with severely reduced kidney function but not on dialysis are not aware of having CKD. Most (96%) people with kidney damage or mildly reduced kidney function are not aware of having CKD”.


The leading causes of the CKD in the united states and most of the world are DM, HTN and Glomerulonephritis.



CKD is defined as decreased kidney function or the presence of kidney damage for three or more months with parameters such as eGFR <60 mL/min/1.73 m² or urinary albumin excretion of ≥30 mg/day¹. In the United States, the prevalence of CKD from 2013-2016 was 14.8 percent (CI=13.6-16.1, with 6.9 percent (CI=6.2-7.6) with stage 3-5 CKD². Medicare spending in 2017 for all patients with CKD was greater than \$84 billion. An additional \$35.9 billion was spent on patients with end stage renal disease (ESRD), representing 7.2% of Medicare’s fee for service spending³. With the rising prevalence and cost of healthcare for patients with CKD and the subsequent progression to ESRD, it is important to identify factors that would lead to earlier detection, better treatment, and identification and control of risk factors.

Morbidity and Mortality of CKD:

The United States Renal Data System (USRDS) reported that in 2016, Medicare patients with CKD had a mortality rate of 122.6 per 1,000 patient years compared to 43.1 for patients without CKD⁴. In addition, rates of all cause hospitalization for CKD increased with disease severity⁴.



In an article published by the International Society of Nephrology, a GFR <60 ml/min/1.73 m² was found to be associated with a greater risk of complications such as drug toxicity, metabolic and endocrine dysfunction, CVD, and death compared to patients with a history of CKD but conserved GFR⁵. There was a relatively constant risk of all outcomes for patients with an eGFR 75-105 ml/min/1.73 m² with an increased relative risk for eGFR of <60 ml/min/1.73 m². This suggests that early detection of renal dysfunction to prevent or slow progression of renal dysfunction can improve morbidity and mortality. They state that due to the high prevalence and cost associated with CKD and its complications, certain countries have developed public health programs aimed towards early detection and treatment. However, follow up studies have yet to determine the efficacy of these measures.

Assessing for microalbuminuria (albumin excretion that exceeds normal physiologic excretion but is below the level of detection for total protein) using albumin-specific dipsticks has been recommended by the National Kidney Foundation for Diabetic patients with increased risk of CKD⁶. If positive, an albumin-creatinine ratio should be obtained to confirm proteinuria within 3 months. A value greater than or equal to 30 mg/g warrants further evaluation. A study showed that screening patients with hypertension, diabetes mellitus, and age greater than 55 yielded a low number needed to screen of 8.7 (95% CI=8.5-9.0)⁷. In subsequent follow up, the risk of end stage renal disease was low, which could suggest decreased morbidity secondary to early detection and increased efforts to slow progression of disease.

Controlling risk factors for CKD is another avenue to consider. Diabetes and hypertension were both shown as independent risk factors that reduce kidney function, and in patients with both risk factors, eGFR was significantly reduced by 25.6%⁸. Diabetic kidney disease was also found to be independently associated with high atherogenic load and increased systolic blood pressure in patients with type 1 diabetes⁹, which not only worsen kidney function but also contribute to the develop of cardiovascular disease. Interestingly, diabetes seems to impact cardiovascular mortality more than CKD mortality. The PREVEND study was a cross sectional cohort study that showed that patients who screened positive for microalbuminuria had the highest incidence of cardiovascular death over a 3 year follow up period¹⁰. Therefore, controlling cardiovascular risk factors in CKD will improve mortality.

Approach and Recommendations for Care

The overall recommendation for early screening of CKD are in high risk patients, such as those with hypertension and diabetes mellitus. Screening with albumin specific dipstick to asses for microalbuminuria has been shown to be the best screening tool with follow up confirmatory albumin to creatinine ratio. In patients with positive screening tests, management of vascular risk factors and comorbidities should adhere to a primary care model¹¹. The American Society of Nephrology suggests an optimal time of consultation for specialists is at stages 3-4¹². Otherwise, lifestyle modifications and pharmacologic interventions for risk factor control can be managed by primary caregivers. These interventions include smoking cessation, dietary salt restriction of less than 2 g daily, protein daily intake restriction to 0.8g/kg of ideal body weight, weight loss to BMI between 20-25, and physical activity of 30 minutes daily for cardiovascular health¹³. In patients with an albumin to creatinine ratio greater than or equal to 30mg/g, blood pressure should be less than 130/80 and started on an ACE inhibitor or ARB¹³. HbA1c should also be maintained at below 7%¹³.³. It is important for physicians, especially those in primary care, to understand the basic principles and steps in screening, managing risk factors, and referring to specialists at an appropriate time. These skills should be emphasized at the medical student level so that they may be prepared to recognize and treat kidney disease appropriately when they begin their training as interns due to the growing prevalence, morbidity, mortality, and healthcare cost associated with CKD.

References

- Saran R, Robinson B, Abbott KC, et al. US Renal Data System 2019 Annual Data Report: epidemiology of kidney disease in the United States. *Am J Kidney Dis.* 2020;75(1)(suppl 1):Svii-Sviii, S1-S64.
- Centers for Disease Control and Prevention. *Chronic Kidney Disease in the United States, 2019.* Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention; 2019. <https://www.cdc.gov/kidneydisease/publications-resources/2019-national-facts.html>
1. Chapter 1: Definition and classification of CKD. *Kidney Int Suppl (2011).* 2013;3(1):19–62. doi:10.1038/kisup.2012.64
 2. Chapter 1: CKD in the General Population. (2018). 2018 USRDS ANNUAL DATA REPORT VOLUME 1: CKD IN THE UNITED STATES. [online] United States Renal Data System, pp. 117-136. Available at:
 3. Chapter 7: Healthcare Expenditures for Persons with CKD. (2018). 2018 USRDS Annual Data Report Volume 1: CKD in the United States. [online] United States Renal Data System, pp. 133-157. Available at: https://www.usrds.org/2018/download/v1_c07_CKD_Costs_18_usrds.pdf.
 4. Chapter 3: Morbidity and Mortality in Patients with CKD. (2018). 2018 USRDS Annual Data Report Volume 1: CKD in the United States. [online] United States Renal Data System, pp.45-78. Available at: https://www.usrds.org/2018/download/v1_c03_MorbMort_18_usrds.pdf.
 5. Chapter 1: Definition and classification of CKD. (2013). *Kidney international supplements*, 3(1), 19–62. doi:10.1038/kisup.2012.64
 6. Levey AS, Coresh J, Balk E, Kausz AT, Levin A, Steffes MW. National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Ann Intern Med* 2003;139:137-47.
 7. Hallan S. Screening strategies for chronic kidney disease in the general population: follow-up of cross sectional health survey. *BMJ.* 2006;334(7585). doi:10.1136/bmj.39098.405382.be.
 8. Bao YS, Jia XB, Ji Y, Yang J, Zhao SL, Na SP. High prevalence and risk factors for kidney dysfunction in patients with atherosclerotic cardio-cerebrovascular disease. *QJM.* 2014;107(6):443-449. doi:10.1093/qjmed/hcu007.
 9. Pacilli, A, Viazzi, F, Fioretto, P, et al. Epidemiology of diabetic kidney disease in adult patients with type 1 diabetes in Italy: The AMD-Annals initiative. *Diabetes Metab Res Rev.* 2017; 33:e2873. doi: 10.1002/dmrr.2873.
 10. Smink PA, Heerspink HJL, Gansevoort RT, et al. Albuminuria, Estimated GFR, Traditional Risk Factors, and Incident Cardiovascular Disease: The PREVEND (Prevention of Renal and Vascular Endstage Disease) Study. *American Journal of Kidney Diseases.* 2012;60(5):804-811. doi:10.1053/j.ajkd.2012.06.017.
 11. Wouters OJ, O'Donoghue DJ, Ritchie J, Kanavos PG, Narva AS. Early chronic kidney disease: diagnosis, management and models of care. *Nat Rev Nephrol.* 2015;11(8):491–502. doi:10.1038/nrneph.2015.85
 12. P Jungers, et al. *Nephrol Dial Transplant* 16: 2357, 2001
 13. Gansevoort RT, Correa-Rotter R, Hemmelgarn BR, et al. Chronic kidney disease and cardiovascular risk: epidemiology, mechanisms, and prevention. *The Lancet.* 2013;382(9889):339-352. doi:10.1016/s0140-6736(13)60595-4.

Biography



Zeidan Hammad, MD is an Assistant Professor of Family Medicine at the Herbert Wertheim College of Medicine in Miami, FL. Dr. Hammad received his medical degree and completed his residency training in Nephrology at the University of Medical Sciences of Camaguey, Cuba. Dr. Hammad also completed a Family Medicine Residency program at the University of Medicine and Dentistry in Hoboken, New Jersey.



Elaine Huang is a fourth year medical student at the Herbert Wertheim College of Medicine in Miami, Florida. She is applying into Internal Medicine.

Article Description:

The prevalence of chronic kidney disease continues to rise globally. Cost, morbidity, and mortality associated with CKD especially in patients 65 years and older are discussed here. Potential avenues of mortality and cost reduction such as early screening, multidisciplinary model of care, and primary care involvement should be considered.